



FOUNDATIONS
of Emergency Medicine

Foundations Frameworks

Approach to the Undifferentiated Rash

Author: Quentin Reuter, MD

Editors: Emilie Fowlkes, MD; Kristen Grabow Moore, MD, MEd

The diagnosis and management of undifferentiated rashes is an expansive and difficult topic. Here is a basic, prioritized way for EM providers to think through the undifferentiated rash. The goal isn't to correctly diagnose every patient's benign, idiopathic rash; the purpose of the ED, as with all medical complaints, is to rule out the life-threatening rashes.

Rash Red Flags: fever, toxic appearance, severe pain, mucosal involvement, new medications, immunosuppressed, very old/young

Can't Miss/immediately Life-Threatening Rashes

1. Purpura/Petechiae: non-blanching, red/purple spots caused by bleeding under the skin

- **Infectious: meningococcemia/septicemia (typically palpable purpura)**
 - Patients are typically febrile, toxic appearing, may have altered mental status
 - Caused by gram negative bacteremia leading to DIC
 - Start broad spectrum antibiotics (cover for meningitis), fluid resuscitate, source control, obtain blood/CSF cultures, ICU admission
- **Microangiopathic hemolytic anemias (MAHA): DIC/HUS/TTP (typically non-palpable purpura)**
 - **Disseminated Intravascular Coagulation (DIC):**
 - Systemic, inappropriate activation of coagulation cascade leading to formation of intravascular fibrin clots, depletion of clotting factors, bleeding, and end organ injury
 - Occurs in setting of sepsis (most common), severe trauma, pregnancy (septic abortion, HELLP), malignancy (leukemia)
 - Low platelets, prolonged hemostasis, low fibrinogen
 - Consider replacing factors if patient bleeding
 - **Hemolytic Uremic Syndrome (HUS):**
 - Mostly seen in children, preceded by diarrheal illness, characterized by hemolytic anemia, thrombocytopenia, acute renal failure (typically with preserved coagulation and fibrinogen)
 - **Thrombotic Thrombocytopenic Purpura (TTP):**
 - TTP is due to destruction of ADAMTS13, a von Willebrand factor multimer cleaving enzyme, which results in abnormal platelet and clot formation
 - Can also be induced by certain medications (quinolones, ticlopidine, clopidogrel), infection, pregnancy
 - Characterized by microangiopathic hemolytic anemia, fever, thrombocytopenia, neurologic abnormalities, and renal disease
- Other causes: **vasculitis, ITP** (afebrile, non-toxic patients)

2. TEN/SJS

- **Stevens-Johnson syndrome (SJS)** and **toxic epidermal necrolysis (TEN)** are severe mucocutaneous reactions, most commonly triggered by medications, characterized by extensive necrosis and detachment of the epidermis.
- Typically begins with ill-defined, coalescing erythematous macules with purpuric centers, can also begin as diffuse erythema, painful in nature
- Progress to vesicles and bullae, positive Nikolsky's sign (slight rubbing of the skin results in exfoliation of the outermost layer)
- Evaluate for mucus membrane involvement (oral, ocular, urogenital)
- Evaluate for recent use of medications that are common precipitants (allopurinol, sulfonyleureas, antibiotics, lamotrigine, NSAIDs)
- Management: removal of inciting exposure, fluids resuscitation, admission to burn unit or ICU

3. Anaphylaxis

- Diffuse hives, erythema/flushing, itching associated with other signs of allergic reaction (nasal/oral swelling, reactive airway disease, hypotension, tachycardia, vomiting and diarrhea)

4. Necrotizing Fasciitis

- Infection of the deeper tissues that results in progressive destruction of the muscle fascia and overlying subcutaneous fat
- Affected area may be erythematous (without sharp margins), swollen, warm, shiny, and *exquisitely tender*, progresses from red-purple to blue-gray patches, then skin breaks down forming bullae and eventually cutaneous gangrene

Other Important Rashes

- **Rocky Mountain Spotted Fever (RMSF)**: Rash develops 2-5 days post onset of fever, rash begins as small, flat, pink, non-itchy spots (macules) on the wrists, forearms, and ankles. Progresses to involve palms and soles, may become petechial rash
- **Lyme**: bull's-eye rash, erythema migrans, forms at site of tick bite, 3 or more days post exposure
- **Bullous pemphigoid**: autoimmune blistering, may be intensely itchy, bullae evolve over weeks to months, start as hives and progress to bullae, Nikolsky's negative
- **Pemphigous vulvaris**: autoimmune blistering, painful but rarely pruritic, commonly begins as oral lesions, + Nikolsky's, prone to secondary infection
- **Staphylococcal scalded skin syndrome (SSSS)**: exotoxin mediating, diffuse erythroderma progress to diffuse areas of exfoliation, Nikolsky's positive, *sparing mucous membranes*, most patients < 2 yo, almost all < 6 yo, admit for fluids, antibiotics, consider transfer to burn center if severe
- **Erythema multiforme**: acute, self-limited, target raised edematous papules distributed peripherally, can involve mucous membranes (major), most caused by HSV (also other viral, bacterial infections, drugs, and RA/SLE), usually self-limited and resolves in 2-6 weeks, punch biopsy to differentiate from SJS/TEN
- **Shingles**: vesicular eruptions along dermatomes, pain/paresthesia/pruritus can precede rash by 2-3 days
- **Scarlet Fever**: caused by group A strep, starts on neck, axillae, groin, spreads to trunk and extremities, red, finely punctate, sandpaper feel, eventually leads to desquamation, treat with antibiotics

- **Kawasaki's Disease:** fever for 5 days plus conjunctivitis, mucus membrane involvement, swelling/peeling in hands/feet, cervical adenopathy, maculopapular rash

Benign Entities

- **Eczema:** erythematous, crusts, fissures, pruritis, excoriations, lichenification, dryness and thickening of skin
- **Contact Dermatitis:** type IV hypersensitivity reaction, itching and burning, exposure to irritant
- **Scabies:** eruption with linear burrows, papules, pustules. Very itchy.

References:

- Nirken, MH., High, WA., Jean-Claude, R. Stevens-Johnson syndrome and toxic epidermal necrolysis: Pathogenesis, clinical manifestations, and diagnosis. uptodate.com
- Kelly, R. Approach to the patient with retiform (angulated) purpura. Apr 2017. uptodate.com
- Campbell, RL, Kelso, JM. Anaphylaxis: Acute diagnosis. July 2016 uptodate.com
- Steven, DL, Baddour LM. Necrotizing soft tissue infections. August 2014. Uptodate.com
- Adams, J et al. Emergency Medicine: Clinical Essentials, Second Edition. Approach to the Adult Rash, pp. 1598-1608. 2013.
- Adams, J et al. Emergency Medicine: Clinical Essentials, Second Edition. Rash in the Severely Ill Patient, pp. 1609-1618. 2013.